

RESEARCH AND PRACTICE

However, we did obtain verbal consent via telephone from each participant before we began the interview process. This process was approved by the Committee on Human Subject Research at the University of California, San Francisco.

References

- Eisenberg DM, Davis RB, Ettner SL, et al. Trends in alternative medicine use in the United States, 1990–1997: results of a follow-up national survey. *JAMA*. 1998;280:1569–1575.
- Lee MM, Lin SS, Wrensch MR, Adler SR, Eisenberg D. Alternative therapies used by women with breast cancer in four ethnic populations. *J Natl Cancer Inst*. 2000;92:42–47.
- Sparber A, Bauer L, Curt G, et al. Use of complementary medicine by adult patients participating in cancer clinical trials. *Oncol Nurs Forum*. 2000;27:623–630.
- Von Gruenigen VE, White LJ, Kirven MS, Showalter AL, Hopkins MP, Jenison EL. A comparison of complementary and alternative medicine use by gynecology and gynecologic oncology patients. *Int J Gynecol Cancer*. 2001;11:205–209.
- Adler SR, Fosket JR. Disclosing complementary and alternative medicine use in the medical encounter: a qualitative study in women with breast cancer. *J Fam Pract*. 1999;48:453–458.
- Nam RK, Fleshner N, Rakovitch E, et al. Prevalence and patterns of the use of complementary therapies among prostate cancer patients: an epidemiological analysis. *J Urol*. 1999;161:1521–1524.
- Kao GD, Devine P. Use of complementary health practices by prostate carcinoma patients undergoing radiation therapy. *Cancer*. 2000;88:615–619.
- Lippert MC, McClain R, Boyd JC, Theodorescu D. Alternative medicine use in patients with localized prostate carcinoma treated with curative intent. *Cancer*. 1999;86:2642–2648.
- Lubeck DP, Litwin MS, Flanders SC, Henning JM, Carroll PR. Use of complementary and alternative medicine (CAM) treatments among a cohort of newly diagnosed prostate cancer patients: data from CAPSUREO [abstract]. *Abstract Book Assoc Health Serv Res*. 1999;16:37.
- Richardson MA, Sanders T, Palmer JL, Greisinger A, Singletary SE. Complementary/alternative medicine use in a comprehensive cancer center and the implications for oncology. *J Clin Oncol*. 2000;18:2505–2514.
- Gotay CC, Hara W, Issell BF, Maskarinec G. Use of complementary and alternative medicine in Hawaii cancer patients. *Hawaii Med J*. 1999;58:94–98.
- Crocetti E, Crotti N, Feltrin A, Ponton P, Geddes M, Buiatti E. The use of complementary therapies by breast cancer patients attending conventional treatment. *Eur J Cancer*. 1998;34:324–328.
- Paltiel O, Avitzour M, Peretz T, et al. Determinants of the use of complementary therapies by patients with cancer. *J Clin Oncol*. 2001;19:2439–2448.
- Leung JM, Dzankic S, Manku K, Yuan S. The prevalence and predictors of the use of alternative medicine in presurgical patients in five California hospitals. *Anesth Analg*. 2001;93:1062–1068.
- Astin JA. Why patients use alternative medicine:

results of a national study. *JAMA*. 1998;279:1548–1553.

- Maskarinec G, Shumay DM, Kakai H, Gotay CC. Ethnic differences in complementary and alternative medicine use among cancer patients. *J Altern Complementary Med*. 2000;6:531–538.
- Smolinske SC. Dietary supplement-drug interactions. *J Am Med Womens Assoc*. 1999;54:191–192, 195.
- Ang-Lee MK, Moss J, Yuan CS. Herbal medicine and perioperative care. *JAMA*. 2001;286:208–216.
- DiPaola RS, Zhang H, Lambert GH, et al. Clinical and biologic activity of an estrogenic herbal combination (PC-SPES) in prostate cancer. *N Engl J Med*. 1998;339:785–791.
- Burstein HJ, Gelber S, Guadagnoli E, Weeks JC. Use of alternative medicine by women with early-stage breast cancer. *N Engl J Med*. 1999;340:1733–1739.

Acupuncture Treatment in the Prevention of Uncomplicated Recurrent Lower Urinary Tract Infections in Adult Women

Terje Alraek, BAC, Liv Inger Fosli Soedal, Siri Urnes Fagerheim, Asbjørn Digranes, MD, and Anders Baerheim, MD, PhD

Acute lower urinary tract infections (UTIs) are common in adult women, and as many as 6% of members of the adult female population experience 3 or more episodes during a given year.¹ In 1995, an estimated 11.3 million women in the United States received antibiotic treatment for at least 1 presumed UTI, resulting in associated costs of \$1.6 billion during that year.² Women with frequently recurrent cystitis may need prophylactic antibacterial treatment. However, such treatment may result in development of antimicrobial resistance, which is a medical problem of increasing concern.³

A recent study indicated that the rate of cystitis among cystitis-prone women treated with acupuncture was one third the rate among untreated women and half the rate

among women treated by sham acupuncture (shallow needling outside known acupuncture points).⁴ In the present study, we sought to evaluate the effect of acupuncture treatment in preventing uncomplicated recurrent lower UTIs among adult nonpregnant women.

METHODS

We recruited 100 women aged 18 to 60 years through advertisements placed in local newspapers in Bergen, Norway. Recruitment took place over a 12-month period beginning in spring 1998. To be included in the study, women had to have had 3 or more episodes of distal urinary symptoms (i.e., dysuria and frequent urination or suprapubic discomfort) during the previous 12 months, and at least 2 of these episodes had to have been diagnosed and treated as acute lower UTIs by a medical doctor. Women were excluded if they were pregnant or were known to have a complicating illness (e.g., diabetes, cancer, obstruction of the urinary tract).

Two women failed to meet the inclusion criteria. Of the 98 women included in the study, 4 subsequently dropped out (2 in the acupuncture group and 2 in the control group).

After written informed consent had been obtained, baseline measurements had been taken, and a diagnosis had been made according to principles of traditional Chinese medicine (TCM), a closed envelope containing the patient's randomized assignment to acupuncture treatment or no treatment was opened. Participants were randomized in blocks of 4 (at a ratio of 3:1) to acupuncture treatment (n=67) or no treatment (n=27).

Treatment consisted of insertion of needles and obtaining of *deqi* (a sensation described as numbness, heaviness, and distention). Acupuncture points were chosen according to the patient's TCM diagnosis. Points were located on the lower abdomen or back (CV-3 or CV-4 and BL-23 or BL-28) or on the lower extremities (KI-3, SP-6, SP-9, ST-36, or LR-3). Treatments were administered twice weekly for 4 weeks.

The primary effect measure was number of occurrences of acute lower UTI during a 6-month follow-up. Residual urine was measured with the automatic Bladder BVI Scan

2500 (Diagnostic Ultrasound Corporation, Wash). Measurements were repeated at 2, 4, and 6 months.

Samples used in bacteriological examinations were collected from clean-voided urine via a dip slide (Uricult; Orion Diagnostica, Finland). Specimens were examined, by bacteriologists unaware of group assignment, at the Gade Institute, Haukeland Hospital, in Bergen, Norway. Acute lower UTI was defined as the presence of (1) the distal urinary symptoms described earlier and (2) bacteriuria (10^5 or more colony-forming units per milliliter of uropathogen or any amount of *Staphylococcus saprophyticus*).

We used Fisher exact tests and χ^2 tests (categorical data) or independent *t* tests (parametric data) to assess differences between groups; differences within groups were examined with paired *t* tests. We report incidence rates of acute lower UTIs in the observation period as number of episodes per person-month, and results of comparisons between groups as incidence rate ratios (IRRs).

RESULTS

As can be seen in Table 1, there were no statistical differences between groups in regard to mean age or mean number of acute lower UTIs in the preceding 12 months. Following treatment, 73% of women in the acupuncture group were free of UTIs during the 6-month observation period, as compared with 52% of women in the control group ($P=.08$). During the observation period, half as many UTI episodes per person-month occurred in the acupuncture group as in the control group (IRR=0.45; 95% confidence interval [CI]=0.23, 0.86; $P\leq.05$; Table 1).

Several episodes of acute distal urinary symptoms were noted in which bacterial cultures either were not obtained or were negative. One third as many episodes per person-month occurred in the acupuncture group as in the control group (IRR=0.30; 95% CI=0.16, 0.58; $P\leq.01$; Table 1).

Women in the acupuncture group experienced a 50% reduction in residual urine after

6 months relative to baseline (35.4 vs 18.2 mL; $P\leq.01$), whereas women in the untreated group exhibited no significant change in residual urine (35.5 vs 38.8 mL). As can be seen in Table 2, the number of women in the treated group with residual urine levels below 10 mL had doubled by the 6-month follow-up, compared with no change among women in the untreated group ($P\leq.05$).

DISCUSSION

Our results showed that acupuncture reduced the recurrence rate among cystitis-prone women to half the rate among untreated women. Also, women in the acupuncture group exhibited reductions in residual urine. The finding that 50% of women in the untreated group became free of cystitis over the study period may reflect regression to the mean, as observed in other studies focusing on recurrent UTIs.^{4,5}

Typically, only 10% of a normal adult female population without specific urinary symptoms will have residual urine levels above 10 mL⁷; 75% of our participants had levels above 10 mL at study entry. Residual urine is a risk factor in the pathogenesis of recurrent lower UTIs among postmenopausal women.^{8,9} Our data indicate that this also may be the case among adult women in general.

In summary, our results, as well as previous findings,⁴ indicate that acupuncture treatment may be effective in preventing recurrent lower UTIs in healthy adult women. ■

TABLE 1—Background Data and UTI Results During a 6-Month Observation Period: 94 Women Receiving Acupuncture Treatment for Recurrent Lower UTIs or Receiving No Treatment

	Acupuncture (n=67)	No Treatment (n=27)
Baseline data		
Mean age, y	38.6	39.1
Mean age at first UTI, y	14.6	14.0
Mean no. of UTIs in previous 12 months	6.2	4.9
Mean no. of weeks since most recent UTI	5.3	6.3
Mean weekly no. of sexual encounters in previous month	1.9	1.7
Main results over 6 months		
Symptomatic lower UTI (with bacteriuria)		
No. of patients without symptoms	49	14
No. of UTIs ^a	18	18
Incidence rate ^b	0.049	0.108
Incidence rate ratio (95% CI)	0.45 (0.23, 0.86)	
UTI symptoms without bacteriuria		
No. of patients without symptoms	52	10
No. of symptomatic episodes	14	21
Incidence rate ^b	0.038	0.125
Incidence rate ratio (95% CI)	0.30 (0.16, 0.58)	

Note. UTI = urinary tract infection. Differences between groups are shown as incidence rate ratios and 95% confidence intervals (CIs).

^aNo participants had more than 2 episodes over the 6-month observation period.

^bCalculated as number of UTIs per person-month.

About the Authors

Terje Alraek and Anders Baerheim are with the Department of Public Health and Primary Health Care, Section for General Practice, University of Bergen, Bergen, Norway. Liv Inger Fosli Soedal and Siri Urnes Fugerheim are medical students in the Department of Public Health and Primary Health Care, Section for General Practice, University of Bergen. Ashjorn Digranes is with the Department of Microbiology and Immunology, Gade Institute, Haukeland Hospital, Bergen, Norway.

Requests for reprints should be sent to Terje Alraek, BAC, Department of Public Health and Primary Health Care, University of Bergen, Ulriksdal 8C, N-5009 Bergen, Norway (e-mail: terje.alraek@isfuib.no).

This brief was accepted June 7, 2002.

Contributors

T. Alraek prepared the study protocol, recruited and treated patients, took part in the data analysis, and

TABLE 2—Residual Urine Levels During a 6-Month Observation Period: 94 Women Receiving Acupuncture Treatment (A) for Recurrent Lower Urinary Tract Infections or Receiving No Treatment (C)

Residual Urine, mL	Baseline, No.		2 Months, No.		4 Months, No.		6 Months, No.	
	A (n=62)	C (n=27)	A (n=64)	C (n=25)	A (n=53)	C (n=18)	A (n=61)	C (n=23)
≤10	15	5	26	9	25	8	32	5*
11-30	20	10	17	7	17	2	14	9
31-100	24	9	14	7	10	7	14	7
>100	3	3	7	2	1	1	1	2

* $P \leq .05$ (vs acupuncture group).

wrote the brief. L. I. F. Soedal and S. U. Fagerheim participated in the data analysis and in preparation of the brief. A. Digranes conducted urinalyses and contributed to the study protocol. A. Baerheim supervised the study and contributed in all phases.

Acknowledgments

This study was funded by the Norwegian Research Council and the Eckbos Legacy.

We thank Norbert Cools, the participating acupuncturist, and Gerd Anne Wincentsen, the nurse who carried out all physical measurements. We also thank Roger Martin for linguistic support.

Human Participant Protection

This study was approved by the Regional Ethical Committee and the Norwegian Data Inspectorate.

References

- Walker M, Heady JA, Shaper AG. The prevalence of dysuria in women in London. *J R Coll Gen Pract*. 1983;33:411-415.
- Foxman B, Barlow R, D'Arcy H, Gillespie B, Sobel JD. Urinary tract infection, self-reported incidence and associated cost. *Ann Epidemiol*. 2000;10:509-515.
- Schaeffer AJ. Urinary tract infections: antimicrobial resistance. *Curr Opin Urol*. 2000;10:23-24.
- Aune A, Alraek T, Huo L, Baerheim A. Acupuncture in the prophylaxis of recurrent lower urinary tract infection in adult women. *Scand J Prim Health Care*. 1998;16:37-39.
- Baerheim A, Larsen E, Digranes A. Vaginal application of lactobacilli in the prophylaxis of recurrent lower urinary tract infection in women. *Scand J Prim Health Care*. 1994;12:239-243.
- O'Grady F, Cattell WR. Kinetics of urinary tract infection, II: the bladder. *Br J Urol*. 1966;38:156-162.
- Raz R, Gennesin Y, Wasser J, et al. Recurrent urinary tract infections in postmenopausal women. *Clin Infect Dis*. 2000;30:152-156.
- Stamm WE, Raz R. Factors contributing to susceptibility of postmenopausal women to recurrent urinary tract infections. *Clin Infect Dis*. 1999;28:723-725.

Yoga for Women With Hyperkyphosis: Results of a Pilot Study

Gail A. Greendale, MD, Anna McDivit, BS, Annie Carpenter, MS, Leanne Seeger, MD, and Mei-Hua Huang, DrPH

The thoracic region of the spine is normally kyphotic, or anteriorly concave. Hyperkyphosis, colloquially called "dowager's hump," refers to excessive kyphotic curvature; however, there is no criterion standard, nor are there any outcome-based definitions of the condition. A kyphosis angle $\geq 40^\circ$ —the 95th percentile value for young adults—is currently used to define hyperkyphosis.^{1,2}

Hyperkyphosis may be associated with physical and emotional limitations³⁻¹¹ and may have multiple precipitants.^{4,12-14} Yoga could be an optimal intervention for hyperkyphosis in that it may improve physical and emotional functioning as well as combat some of the underlying muscular and biomechanical causes. We conducted a single-arm, non-masked intervention trial to assess the effects on anthropometric and physical function of yoga among women with hyperkyphosis.

METHODS

To be included in the study, which was conducted in Los Angeles during September 2000 to September 2001, women had to meet the following criteria: presence of physician-diagnosed hyperkyphosis, age 60 years

or older, absence of angina and uncontrolled lung disease, cleared for participation by primary care physician, and able to pass physical safety tests (e.g., able to rise from the floor to a standing position safely and independently). The intervention involved hatha yoga, a type of yoga incorporating a combination of breathing and movement.

As a means of ensuring the safety of the participants, the study took place in a closely monitored environment involving one-on-one supervision and hands-on adjustments and corrections. The women were divided into 2 separate small classes (n=11 and n=10), each of which involved 12 weeks of yoga consisting of twice-weekly 1-hour sessions.

The program included 4 series of poses modified from the classical forms of yoga to accommodate the physical constraints of kyphotic women. More challenging poses were introduced every 3 weeks, and muscles and joints particularly affected by hyperkyphosis (shoulders, spinal erectors, abdominals, neck) were targeted. Figure 1 briefly summarizes the 4 series and illustrates an example of 1 pose from each.

Anthropometric outcomes, assessed at baseline and follow-up by 1 of the investigators (A.M.) by means of standard protocols, were (1) height without shoes (measured with a stadiometer), (2) distance from tragus to wall (a measure of forward curvature), and (3) Debrunner kyphometer angle (an estimate of degree of thoracic spinal curvature; higher values indicate more curvature).¹⁵ At baseline, this investigator performed same-day repeated measurements of each anthropometric characteristic for 6 of the participants. Intra-class correlation coefficients were .98, .61, and .34, respectively.

Timed physical performance measures were chair stands (standing up and sitting down, with arms folded across chest, using an armless chair),¹⁶ functional reach,¹⁷ the "penny test" (picking up a penny from the floor),¹⁸ the "book test" (placing a book on a high shelf),¹⁸ and an 8-ft (2.4-m) walk.¹⁹ At baseline, all participants underwent spinal radiographs; radiographs were read by a skeletal radiologist.²⁰

We used pretest-posttest scores to compute changes in each anthropometric and performance outcome. Because our sample size